arXiv:1603.03965v1 [math.CA] 12 Mar 2016

ON PALEY-TYPE AND HAUSDORFF – YOUNG – PALEY-TYPE INEQUALITIES FOR JACOBI EXPANSIONS

Roman A. Veprintsev

Abstract. We obtain Paley-type and Hausdorff – Young – Paley-type inequalities for Jacobi expansions.

Key words and phrases: orthogonal polynomials, Jacobi polynomials, Paley-type inequality, Hausdorff – Young – Paley-type inequality

MSC 2010: 33C45, 41A17, 42C10

1. Introduction and preliminaries

In this section, we introduce the basic notation and review some results on Jacobi polynomials. Here, we also formulate the aim of the publication.

Let $\mathbb{N}_0$ denote the set of non-negative integers. Suppose that α, $\beta > -1$. The Jacobi polynomials, denoted by $P_n^{(\alpha,\beta)}(\cdot)$, $n \in \mathbb{N}_0$, are orthogonal with respect to the Jacobi weight function $w_{\alpha,\beta}(t) = (1-t)^\alpha (1+t)^\beta$ on $[-1,1]$, namely,

$$\int_{-1}^{1} P_n^{(\alpha,\beta)}(t)\, P_m^{(\alpha,\beta)}(t)\, w_{\alpha,\beta}(t)\, dt = \begin{cases} \dfrac{2^{\alpha+\beta+1}\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{(2n+\alpha+\beta+1)\Gamma(n+1)\Gamma(n+\alpha+\beta+1)}, & n=m, \\ 0, & n \neq m. \end{cases}$$

Here, as usual, Γ is the gamma function.

We use the following asymptotic notation: $f(n) \asymp g(n)$, $n \to \infty$, means that there exist positive constants C_1, C_2, and a positive integer n_0 such that $0 \le C_1 g(n) \le f(n) \le C_2 g(n)$ for all $n \ge n_0$. To simplify the writing, we will omit "$n \to \infty$" in the notation.

It follows directly from Stirling's asymptotic formula that $\frac{\Gamma(n+\lambda)}{\Gamma(n+\mu)} \asymp n^{\lambda-\mu}$ for arbitrary real numbers λ and μ. Thus, we have

$$h_n^{(\alpha,\beta)} = \Big(\int_{-1}^{1} \big(P_n^{(\alpha,\beta)}(t)\big)^2\, w_{\alpha,\beta}(t)\, dt \Big)^{1/2} \asymp n^{-1/2}. \tag{1}$$

Define the uniform norm of a continuous function f on $[-1,1]$ by

$$\|f\|_\infty = \max_{-1\le t\le 1} |f(t)|.$$

The maximum of two real numbers x and y is denoted by $\max(x,y)$.

It is known [4, Theorem 7.32.1] that

$$\big\| P_n^{(\alpha,\beta)} \big\|_\infty \asymp n^{\sigma(\alpha,\beta)-\frac{1}{2}}, \qquad \text{where} \quad \sigma(\alpha,\beta) = \max\Big(0, \frac{1}{2} + \max(\alpha,\beta)\Big). \tag{2}$$

Let $\big\{ \widetilde{P}_n^{(\alpha,\beta)} \big\}_{n=0}^{\infty}$ be the sequence of orthonormal Jacobi polynomials (with respect to the Jacobi weight function $w_{\alpha,\beta}$), that is, $\widetilde{P}_n^{(\alpha,\beta)} = \big(h_n^{(\alpha,\beta)}\big)^{-1} P_n^{(\alpha,\beta)}$, $n \in \mathbb{N}_0$.

It follows from (1) and (2) that

$$\big\|\widetilde{P}_n^{(\alpha,\beta)}\big\|_\infty \asymp n^{\sigma(\alpha,\beta)}. \tag{3}$$

Given $1 \le p \le \infty$, we denote by $L_p(w_{\alpha,\beta})$ the space of complex-valued Lebesgue measurable functions f on $[-1,1]$ with finite norm

$$\|f\|_{L_p(w_{\alpha,\beta})} = \Big(\int_{-1}^1 |f(t)|^p\, w_{\alpha,\beta}(t)\,dt\Big)^{1/p}, \qquad 1 \le p < \infty,$$

$$\|f\|_{L_\infty} = \operatorname*{ess\,sup}_{x\in[-1,1]} |f(x)|, \qquad p = \infty.$$

For a function $f \in L_p(w_{\alpha,\beta})$, $1 \le p \le \infty$, the Jacobi expansion is given by

$$f(t) \sim \sum_{n=0}^\infty \hat{f}_n \widetilde{P}_n^{(\alpha,\beta)}(t), \qquad \text{where} \quad \hat{f}_n = \int_{-1}^1 f(t)\,\widetilde{P}_n^{(\alpha,\beta)}(t)\,w_{\alpha,\beta}(t)\,dt.$$

For $1 \le p \le \infty$, we denote by p' the conjugate exponent to p, that is, $\frac{1}{p} + \frac{1}{p'} = 1$.

Our aim is to establish Paley-type and Hausdorff – Young – Paley-type inequalities for Jacobi expansions in Sections 2 and 3, respectively.

2. Paley-type inequalities for Jacobi expansions

We will omit the proof of the following Paley-type inequalities for Jacobi expansions because its proof is analogous to that of Theorem 2 given in [5, Section 2].

Theorem 1. *(a) If $1 < p \le 2$, $f \in L_p(w_{\alpha,\beta})$, ω is a positive function on $\mathbb{N}_0$ such that*

$$M_\omega = \sup_{t>0}\Big\{t\Big(\sum_{\omega(n)\ge t} (n+1)^{2\sigma(\alpha,\beta)}\Big)\Big\} < \infty, \tag{4}$$

then

$$\Big\{\sum_{n=0}^\infty \Big((n+1)^{\left(\frac{1}{p}-\frac{1}{p'}\right)\sigma(\alpha,\beta)}\,(\omega(n))^{\frac{1}{p}-\frac{1}{p'}}|\hat{f}_n|\Big)^p\Big\}^{1/p} \le A_p M_\omega^{\frac{1}{p}-\frac{1}{p'}}\|f\|_{L_p(w_{\alpha,\beta})}. \tag{5}$$

(b) If $2 \le q < \infty$, ω is a positive function on $\mathbb{N}_0$ satisfying (4) and ϕ is a non-negative function on $\mathbb{N}_0$ such that

$$\sum_{n=0}^\infty \Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\sigma(\alpha,\beta)}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}|\phi(n)|\Big)^q < \infty,$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^N \phi(n)\widetilde{P}_n^{(\alpha,\beta)}(t)$$

converge in $L_q(w_{\alpha,\beta})$ to a function f satisfying $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, and

$$\|f\|_{L_q(w_{\alpha,\beta})} \le A_{q'} M_\omega^{\frac{1}{q'}-\frac{1}{q}}\Big\{\sum_{n=0}^\infty \Big((n+1)^{\left(\frac{1}{q}-\frac{1}{q'}\right)\sigma(\alpha,\beta)}(\omega(n))^{\frac{1}{q}-\frac{1}{q'}}|\phi(n)|\Big)^q\Big\}^{1/q}.$$

3. Hausdorff – Young – Paley-type inequalities for Jacobi expansions

In [2, Theorem 3.1], Z. Ditzian proved the following Hausdorff – Young-type inequalities for Jacobi expansions.

Theorem 2. (*a*) *If* $1 < p \le 2$ *and* $f \in L_p(w_{\alpha,\beta})$, *then*

$$\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{p'}-\frac{1}{p}\right)\sigma(\alpha,\beta)}|\hat{f}_n|\Big)^{p'}\Big\}^{1/p'} \le B_p\,\|f\|_{L_p(w_{\alpha,\beta})}.$$

(*b*) *If* $2 \le q < \infty$ *and* ϕ *is a function on non-negative integers satisfying*

$$\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\sigma(\alpha,\beta)}|\phi(n)|\Big)^{q'} < \infty,$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^{N}\phi(n)\,\widetilde{P}_n^{(\alpha,\beta)}(t)$$

converge in $L_q(w_{\alpha,\beta})$ *to a function* f *satisfying* $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, *and*

$$\|f\|_{L_q(w_{\alpha,\beta})} \le B_{q'}\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{1}{q'}-\frac{1}{q}\right)\sigma(\alpha,\beta)}|\phi(n)|\Big)^{q'}\Big\}^{1/q'}.$$

Theorem 3 contains the Paley-type and the Hausdorff – Young-type inequalities for Jacobi expansions. The proof of this theorem is analogous to the proof of Theorem 5 in [5, Section 3] and we shall omit it.

Theorem 3. (*a*) *If* $1 < p \le 2$, $f \in L_p(w_{\alpha,\beta})$, ω *is a positive function on* $\mathbb{N}_0$ *satisfying the condition* (4), *and* $p \le s \le p'$, *then*

$$\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(\frac{2}{s}-1\right)\sigma(\alpha,\beta)}(\omega(n))^{\frac{1}{s}-\frac{1}{p'}}|\hat{f}_n|\Big)^{s}\Big\}^{1/s} \le C_p(s)\,M_\omega^{\frac{1}{s}-\frac{1}{p'}}\,\|f\|_{L_p(w_{\alpha,\beta})}.$$

(*b*) *If* $2 \le q < \infty$, $q' \le r \le q$, ω *is a positive function on* $\mathbb{N}_0$ *satisfying the condition* (4) *and* ϕ *is a non-negative function on* $\mathbb{N}_0$ *such that*

$$\sum_{n=0}^{\infty}\Big((n+1)^{\left(1-\frac{2}{r}\right)\sigma(\alpha,\beta)}(\omega(n))^{\frac{1}{q}-\frac{1}{r}}|\phi(n)|\Big)^{r'} < \infty,$$

then the algebraic polynomials

$$\Phi_N(t) = \sum_{n=0}^{N}\phi(n)\widetilde{P}_n^{(\alpha,\beta)}(t)$$

converge in $L_q(w_{\alpha,\beta})$ *to a function* f *satisfying* $\hat{f}_n = \phi(n)$, $n \in \mathbb{N}_0$, *and*

$$\|f\|_{L_q(w_{\alpha,\beta})} \le C_{q'}(r)M_\omega^{\frac{1}{r}-\frac{1}{q}}\Big\{\sum_{n=0}^{\infty}\Big((n+1)^{\left(1-\frac{2}{r}\right)\sigma(\alpha,\beta)}(\omega(n))^{\frac{1}{q}-\frac{1}{r}}|\phi(n)|\Big)^{r'}\Big\}^{1/r'}.$$

4. Remarks

In the following theorem we show that the inequality (5) does not hold for $p = 1$ in general.

Theorem 4. *Under the conditions and notations of* Theorem 1, (5) *is not satisfied for* $p = 1$ *provided that* $\max(\alpha, \beta) \geq -\frac{1}{2}$ *and*

$$\sum_{n=0}^{\infty} \omega(n)(n+1)^{2\sigma(\alpha,\beta)} = +\infty.$$

P r o o f. We assume without loss of generality that $\max(\alpha, \beta) = \alpha$. Then we have $\sigma(\alpha, \beta) = \frac{1}{2} + \alpha$. We define the sequence of polynomials g_N by

$$g_N(t) = \sum_{n=0}^{N} \omega(n)(n+1)^{\sigma(\alpha,\beta)} \widetilde{P}_n^{(\alpha,\beta)}(t).$$

As $\big\|\widetilde{P}_n^{(\alpha,\beta)}\big\|_\infty = \widetilde{P}_n^{(\alpha,\beta)}(1) \asymp (n+1)^{\sigma(\alpha,\beta)}$ (see [4, Theorem 7.32.1, (4.1.1)] and (3)), we obtain $\|g_N\|_\infty = \sum\limits_{n=0}^{N} \omega(n)(n+1)^{\sigma(\alpha,\beta)} \widetilde{P}_n^{(\alpha,\beta)}(1) \asymp \sum\limits_{n=0}^{N} \omega(n)(n+1)^{2\sigma(\alpha,\beta)}$. Thus,

$$\|g_N\|_\infty \to \infty \quad \text{as} \quad N \to \infty.$$

We now show that assuming (5) for $p = 1$ would lead to a contradiction with what we already proved. Because of $\|g_N\|_\infty = \sup\big\{\int_{-1}^{1} g_N(t) f(t)\, w_{\alpha,\beta}(t)\, dt\colon\ \|f\|_{L_1(w_{\alpha,\beta})} = 1\big\}$, we have

$$\begin{aligned}
\|g_N\|_\infty &= \sup\left\{\int_{-1}^{1} g_N(t) f(t)\, w_{\alpha,\beta}(t)\, dt\colon\ \|f\|_{L_1(w_{\alpha,\beta})} = 1\right\} = \\
&= \sup\left\{\sum_{n=0}^{N} \omega(n)(n+1)^{\sigma(\alpha,\beta)} \hat{f}_n\colon\ \|f\|_{L_1(w_{\alpha,\beta})} = 1\right\} \leq \\
&\leq \sup\left\{\sum_{n=0}^{N} \omega(n)(n+1)^{\sigma(\alpha,\beta)} |\hat{f}_n|\colon\ \|f\|_{L_1(w_{\alpha,\beta})} = 1\right\} \leq A_1 M_\omega.
\end{aligned}$$

This is a contradiction, which proves that our assumption is wrong. □

5. Conclusion

As an application of the results above, we are going to obtain sufficient condition for a special sequence of positive real numbers to be a Fourier multiplier.

Acknowledgements

The author would like to thank Michael Ruzhansky for the idea of this work (see the e-print [1]). This work was done thanks to the remarkable paper [2] (see especially Section 4 there) of Z. Ditzian.

REFERENCES

1. R. Akylzhanov, E. Nursultanov, and M. Ruzhansky, Hardy – Littlewood, Hausdorff – Young – Paley inequalities, and L^p-L^q Fourier multipliers on compact homogeneous manifolds, arXiv preprint 1504.07043 (2015).
2. Z. Ditzian, Norm and smoothness of a function related to the coefficients of its expansion, *J. Approx. Theory* **196** (2015), 101–110.
3. E. Stein, Interpolation of linear operators, *Trans. Amer. Math. Soc.* **83** (1956), 482–492.
4. G. Szegö, *Orthogonal polynomials*, 4th ed., American Mathematical Society Colloquium Publications **23**, American Mathematical Society, Providence, Rhode Island, 1975.
5. R. A. Veprintsev, On Paley-type and Hausdorff – Young – Paley-type inequalities for generalized Gegenbauer expansions, arXiv preprint 1603.02072 (2016).

Independent researcher, Uzlovaya, Russia
E-mail address: **veprintsevroma@gmail.com**